



Synthesis of 5-(trifluoromethyl)-5H-chromeno[3,4-b]pyridines from 3-nitro-2-(trifluoromethyl)-2H-chromenes and aminoenones derived from acetylacetone and cyclic amines

Vladislav Yu. Korotaev, Alexey Yu. Barkov, Vyacheslav Ya. Sosnovskikh *

Department of Chemistry, Ural Federal University, pr. Lenina 51, 620000 Ekaterinburg, Russia

ARTICLE INFO

Article history:

Received 15 February 2013

Revised 20 March 2013

Accepted 28 March 2013

Available online 11 April 2013

Keywords:

3-Nitro-2H-chromenes

Push–pull enamines

Michael addition

Chromanes

Chromeno[3,4-b]pyridines

ABSTRACT

Reactions of 3-nitro-2H-chromenes with aminoenones derived from acetylacetone and cyclic amines proceed diastereoselectively to give functionalized 2,3,4-trisubstituted chromanes as a result of nucleophilic addition of the vinylogous β -methyl group at the C-4 atom of the chromene system. From these compounds, under acidic conditions, 5-(trifluoromethyl)-5H-chromeno[3,4-b]pyridines and 4-acetoacetyl-3-nitro-2-(trichloromethyl/phenyl)chromanes, depending on the substituent at the 2-position, were obtained in moderate to good yields.

© 2013 Elsevier Ltd. All rights reserved.

Many derivatives of chromane (3,4-dihydro-2H-1-benzopyran) and 2H-chromene (2H-1-benzopyran) are natural compounds that are widely abundant in the plant kingdom.¹ Some of these, as well as a series of synthetic 2H-chromenes, have found use as pesticides and promising drugs.² 3-Nitro-2H-chromenes, as conjugated nitro-olefins, possess unique chemical reactivity in both nucleophilic and cycloaddition reactions because of their reactive double bond. As a result, these compounds have attracted attention as excellent building blocks for the preparation of various more complex heterocyclic compounds.³ In particular, 3-nitro-2H-chromenes react with diverse C-, N-, S-, and P-nucleophiles to give a wide range of substituted⁴ and fused⁵ chromanes. Recently, the reactions of 2-aryl-3-nitro-2H-chromenes with α,α -dicyanoolefins in the presence of triethylamine, leading to the preparation of 6-aryl-6H-dibenzo[b,d]pyran derivatives, were reported.⁶ At the same time, to the best of our knowledge, very little information is available on the reactions of 3-nitro-2H-chromenes with push–pull enamines.⁷ It has been reported that 2-aryl-3-nitro-2H-chromenes react with methyl β -methylaminocrotonate to give mixtures of the addition product, methyl 3-methylamino-2-(2-aryl-3-nitrochroman-4-yl)-2-butenate, and 1-benzopyrano[3,4-b]pyrrole, formed via the Grob cyclization,⁸ whereas their reactions with ethyl β -morpholinocrotonate took an entirely different course and gave ethyl 3-morpholino-4-(2-aryl-3-nitrochroman-4-yl)-2-butenates.⁹ The


latter result belongs to a rare case, where the reaction proceeded not at the more nucleophilic α -position of an enamino ester, but at the vinylogous β -methyl group. This is of particular interest because it results in the formation of a functionalized chromane system, which has not been reported previously.

Within the framework of a research program on the synthetic opportunities offered by 3-nitro-2-trihalomethyl-2H-chromenes **1** for the preparation of organic molecules of potential interest in biomedical chemistry and materials science,⁵ we examined the behavior of compounds **1** in reactions with the push–pull enamines derived from acetylacetone and cyclic amines, namely (*E*)-4-morpholino- and (*E*)-4-piperidinopent-3-en-2-ones, with the purpose of studying the structures and acid-catalyzed transformations of the addition products **2**. The present communication describes a new type of pyridine ring annulation, which consists of the conversion of compounds **2** into 5-(trifluoromethyl)-5H-chromeno[3,4-b]pyridines **4**, and demonstrates the dramatic effect of the trifluoromethyl group on the reaction pathway.

We found that the reaction of chromenes **1** with 4-morpholino- and 4-piperidinopent-3-en-2-ones in dry acetonitrile at room temperature for 2–48 h resulted in the stereoselective formation of 2,3,4-trisubstituted chromanes **2a–i** in 42%–83% yields as single diastereomeric products with *cis*–*trans*-configuration ($^3J_{H2,H3} = ^3J_{H3,H4} = 1.5$ Hz) at the C(2)–C(3) and C(3)–C(4) bonds, respectively, and with *E*-configuration at the double bond (Scheme 1, Table 1).¹⁰ The stereochemistry of the products was confirmed by an X-ray diffraction study of crystals of **2a** (Fig. 1).¹¹

* Corresponding author. Fax: +7 343 261 59 78.

E-mail address: vyacheslav.sosnovskikh@usu.ru (V.Y. Sosnovskikh).



Scheme 1. Synthesis of compounds 2–4.

Table 1
Synthesis of compounds 2–4

R	R'	X	Product 2	Time (h)	Yield (%)	Product 3, 4	Yield (%)
CF ₃	H	CH ₂	2a	2	71	4a	34
CF ₃	H	O	2b	5	83	4a	33
CF ₃	Br	O	2c	3	80	4c	46
CF ₃	NO ₂	O	2d	6	70	4d	42
CF ₃	MeO	O	2e	48	64	4e	34
CCl ₃	H	O	2f	30	42	3f^a	56
CCl ₃	Br	CH ₂	2g	7	56	3g^b	63
CCl ₃	Br	O	2h	21	52	3g	66
Ph	H	O	2i	7	62	3i^c	60

^a **3f**:**3f'** = 88:12.^b **3g**:**3g'** = 91:9.^c **3i**:**3i'** = 69:31.

The reaction time varied according to the nature of the chromene molecule. In general, 2-CF₃-chromenes were more reactive than 2-CCl₃- and 2-Ph-chromenes; conjugate addition of an enamine to a CF₃-chromene containing an electron-donating MeO group required a longer period of time for completion of the reaction, while electron-withdrawing groups (Br, NO₂) required relatively short reaction times. Addition of push–pull enamines at the vinyl-



ous β-Me group possibly results from the *E*-configuration of the starting enamines, which hinders the approach of the α-C atom to the activated double bond of the chromene system.

Next we examined the acid hydrolysis of enamino ketones **2a-i** in aqueous ethanol at reflux in the presence of concentrated HCl. As expected, hydrolysis of compounds **2f-i** successfully removed the amino function to give interesting acetoacetyl derivatives **3f,g**, and **i** as a mixture of keto-enol tautomers **3** and **3'** in 56–66% yields, with the same configuration (9%–31% of diketo form **3'**). As for the 2-CF₃-chromenes **2a-e**, there was a marked difference in their reactivity toward acid hydrolysis. Under the same conditions, instead of the corresponding acetoacetyl derivatives **3**, we obtained 5-(trifluoromethyl)-5*H*-chromeno[3,4-*b*]pyridines **4a,c-e** in 33%–46% isolated yields (Scheme 1, Table 1). The structures of compounds **3** and **4** were established by IR, ¹H, ¹⁹F, and ¹³C NMR spectroscopy as well as by elemental analysis.¹²

These results indicate that the nature of the substituent at the 2-position influences the course of the acid-catalyzed hydrolysis of **2**, which in the case of a CF₃ group took an unexpected direction compared with the reaction of nonfluorinated chromenes. It is important to note that this new cyclization process tolerates both electron-donating (MeO) and electron-withdrawing (Br, NO₂) substituents on the benzene ring. Moreover, unlike previously known approaches for the synthesis of quinolines from nitro derivatives,¹³ this reaction does not require any reducing agent or chromatographic purification of the final products, and thereby greatly facilitates the preparation of the target chromenopyridines **4**.

We can assume that cyclization of **2** into **4** was possible due to the high C–H acidity of the H-2 in *aci*-form **A**, which could be easily abstracted by the base to form **B** (nitron–*N*-hydroxyenamine prototropic tautomeric equilibria through a [1,4]-H shift¹⁴), followed by dehydration of intermediate **B** into the corresponding α-nitrosoalkene **C**. Intramolecular nucleophilic attack of the side-chain enol on the nitroso group followed by dehydration results in the intermediate **D**, which undergoes a double [1,3]-H shift to give chromenopyridines **4** (Scheme 2). The lack of such reactivity in the case of compounds **2f-i** was apparently due to the lower acidity of the H-2 proton. It should be noted that a similar [1,4]-H shift has been previously observed by us in the spontaneous ring-contraction–rearrangement of CF₃-containing 1,2-oxazine *N*-oxides into 1-pyrroline *N*-oxides.¹⁵

The main information for the characterization of compounds **4** was obtained from the ¹H NMR spectra, which showed no methylene signal, a result consistent with the pyridine structure. The most downfield shifted signal was assigned to the hydroxyl proton, which appeared as a singlet at δ 12.0 in CDCl₃, the singlet

Figure 1. X-ray crystal structure of *cis-trans*-**2a** (ORTEP drawing, 50% probability level).Scheme 2. Mechanism of the formation of chromeno[3,4-*b*]pyridines **4**.

at δ 7.58–7.77 was due to the resonance of the pyridine H-1 proton; assignment of all the signals was achieved by 2D ^1H – ^{13}C HSQC, and HMBC experiments. In the 2D ^1H – ^{15}N HMBC spectrum of **4c**, a signal was observed at δ 316.9 from liquid NH_3 , indicating that the N was present in a pyridine ring.¹⁶ In addition, the CF_3 group in the ^{19}F NMR spectra of chromenopyridines **4** manifested itself as a doublet at δ 85.0–85.5 ($J = 7.0$ – 7.3 Hz).

In conclusion, we have shown, for the first time, that the reaction of 3-nitro-2H-chromenes with push–pull enamines affords linear enamines with the chromanyl moiety, acid hydrolysis of which gave either 5H-chromeno[3,4-b]pyridines or 4-(acetoacetyl)chromanes, depending on the nature of the substituent at C-2. When this substituent was a CF_3 group, a novel synthesis of fused pyridines followed, while acid hydrolysis of CCl_3 - and Ph-containing derivatives proceeded in the expected manner to give interesting polyfunctional chromane derivatives. Further studies on the synthetic scope of these reactions are now in progress.

Acknowledgment

This work was supported financially by the Russian Foundation for Basic Research (Grant 11-03-00126).

References and notes

- (a) Schweizer, E. E.; Meeder-Nycz, D. In *The Chemistry of Heterocyclic Compounds: Chromenes, Chromanones, and Chromones*; Ellis, G. P., Ed.; Wiley: New York, 1977; Vol. 31; (b) *The Chemistry of Heterocyclic Compounds: Chromans and Tocopherols*; Ellis, G. P., Lockhard, I. M., Eds.; Wiley: New York, 1981; Vol. 36; (c) Horton D. A.; Bourne, G. T.; Smythe, M. L. *Chem. Rev.* **2003**, *103*, 893.
- (a) Bowers, W. S.; Ohta, T.; Cleere, J. S.; Marsella, P. A. *Science* **1976**, *193*, 542; (b) Bergmann, R.; Gericke, R. J. *Med. Chem.* **1990**, *33*, 492; (c) Burrell, G.; Cassidy, F.; Evans, J. M.; Lightowler, D.; Stemp, G. J. *Med. Chem.* **1990**, *33*, 3023; (d) Gericke, R.; Harting, J.; Lues, I.; Schittenhelm, C. J. *Med. Chem.* **1991**, *34*, 3074.
- (a) Ono, N. *The Nitro Group in Organic Synthesis*; Wiley-VCH: New York, 2001; (b) Yoshikoshi, A.; Miyashita, M. *Acc. Chem. Res.* **1985**, *18*, 284; (c) Bhanja, C.; Jena, S.; Nayak, S.; Mohapatra, S. *Beilstein J. Org. Chem.* **2012**, *8*, 1668.
- (a) Korotaev, V. Y.; Sosnovskikh, V. Y.; Kutyshev, I. B.; Kodess, M. I. *Lett. Org. Chem.* **2005**, *2*, 616; (b) Korotaev, V. Y.; Sosnovskikh, V. Y.; Kutyshev, I. B.; Kodess, M. I. *Izv. Akad. Nauk, Ser. Khim.* **2006**, 309 (Russ. Chem. Bull., Int. Ed. 2006, *55*, 317); (c) Korotaev, V. Y.; Sosnovskikh, V. Y.; Kutyshev, I. B.; Kodess, M. I. *Izv. Akad. Nauk, Ser. Khim.* **2006**, 1945 (Russ. Chem. Bull., Int. Ed. 2006, *55*, 2020); (d) Korotaev, V. Y.; Kutyshev, I. B.; Sosnovskikh, V. Y.; Kodess, M. I. *Mendeleev Commun.* **2007**, *17*, 52; (e) Korotaev, V. Y.; Sosnovskikh, V. Y.; Kutyshev, I. B. *Izv. Akad. Nauk, Ser. Khim.* **2007**, 1985 (Russ. Chem. Bull., Int. Ed. 2007, *56*, 2054); (f) Biswas, S.; Maulik, P. R.; Gupta, R. C.; Seth, M.; Bhaduri, A. P. *Acta Crystallogr., Sect. C* **1996**, *52*, 1036; (g) Lin, C.; Hsu, J.; Sastry, M. N. V.; Fang, H.; Tu, Z.; Liu, J.-T.; Yao, C.-F. *Tetrahedron* **2005**, *61*, 11751; (h) Ono, N.; Banshou, N.; Ioto, S.; Murashima, T.; Ogawa, T. J. *Heterocycl. Chem.* **1997**, *34*, 1243.
- (a) Korotaev, V. Y.; Sosnovskikh, V. Y.; Barabanov, M. A.; Yasnova, E. S.; Ezhikova, M. A.; Kodess, M. I.; Slepukhin, P. A. *Tetrahedron* **2010**, *66*, 1404; (b) Korotaev, V. Y.; Sosnovskikh, V. Y.; Kutyshev, I. B.; Barkov, A. Y.; Shklyayev, Y. V. *Tetrahedron Lett.* **2008**, *49*, 5376; (c) Korotaev, V. Y.; Sosnovskikh, V. Y.; Barkov, A. Y.; Slepukhin, P. A.; Ezhikova, M. A.; Kodess, M. I.; Shklyayev, Y. V. *Tetrahedron* **2011**, *67*, 8685.
- (a) Li, P.; Luo, L.-L.; Li, X.-S.; Xie, J.-W. *Tetrahedron* **2010**, *66*, 7590; (b) Guo, Z.-W.; Li, X.-S.; Zhu, W.-D.; Xie, J.-W. *Eur. J. Org. Chem.* **2012**, 6924.
- Kostyuk, A. N.; Volochnyuk, D. M.; Sibgatulin, D. A. *Synthesis* **2008**, 161.
- Gupta, R. C.; Seth, M.; Bhaduri, A. P. *J. Indian Chem.* **1991**, *30B*, 297.
- Korotaev, V. Y.; Barkov, A. Y.; Sokovnina, A. A.; Sosnovskikh, V. Y. *Mendeleev Commun.* **2013**, *23*, N 3.
- General procedure for the synthesis of chromanes **2**. A mixture of chromene **1** (1.0 mmol) and (E)-4-morpholino(piperidino)pent-3-en-2-one (0.17 g, 1.0 mmol) in dry MeCN was stirred at room temperature for several hours (Table 1). The solid that formed was filtered and recrystallized from CH_2Cl_2 /hexane (1:2) to give compound **2** as a white powder.
(E)-5-[(2R*,3R*,4S*)-4-Morpholino-3-nitro-2-(trifluoromethyl)-3,4-dihydro-2H-chromen-4-yl]-3-penten-2-one (**2b**). Yield 0.35 g (83%), mp 164–165 °C (dec.). IR (KBr): 1655, 1632, 1549, 1490, 1448, 1374 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 2.13 (s, 3H, Me), 2.75 (dd, 1H, CHH, $J = 14.6$, 2.6 Hz), 3.30 (dt, 2H, N(CHH)₂, $J = 12.9$, 4.9 Hz), 3.30–3.40 (m, 3H, H-4, N(CHH)₂), 3.76 (t, 4H, O(CH₂)₂, $J = 4.9$ Hz), 4.23 (br t, 1H, CHH, $J = 13.3$ Hz), 5.16 (br s, 1H, H-3), 5.27 (br q, 1H, H-2, $J = 6.0$ Hz), 5.47 (s, 1H, =CH), 7.02–7.10 (m, 2H, H-6, H-8), 7.17 (br d, 1H, H-5, $J = 7.3$ Hz), 7.22–7.30 (m, 1H, H-7); ^{13}C NMR (126 MHz, CDCl_3) δ 32.0, 32.5, 38.6, 47.0, 66.2, 70.7 (q, C-2, $^2J_{\text{CF}} = 34.1$ Hz), 77.6, 101.8, 117.6, 120.7, 122.3 (q, CF_3 , $^1J_{\text{CF}} = 280.8$ Hz), 122.8, 128.3, 128.9, 152.0, 158.3, 196.5 (C=O); ^{19}F NMR (376 MHz, CDCl_3) δ 86.8 (d, CF_3 , $J = 6.0$ Hz). Anal. calcd for $\text{C}_{19}\text{H}_{21}\text{F}_3\text{N}_2\text{O}_5$: C, 55.07; H, 5.11; N, 6.76. Found: C, 54.90; H, 4.97; N, 6.82.
- X-ray diffraction study of compound **2a**. Diffraction data were collected at 295 K on an Xcalibur 3 automatic single-crystal diffractometer (graphite-monochromated MoK α radiation, ω -scans). The structure was solved by direct method and refined by the full-matrix least-squares method using the SHELX-97 program package.¹⁷ The H atoms were located geometrically using the riding model. Crystal data for **2a**: $\text{C}_{20}\text{H}_{23}\text{F}_3\text{N}_2\text{O}_4$, $M = 412.40$, monoclinic crystals, space group C2/c , $a = 19.979(2)$, $b = 10.1081(8)$, $c = 20.1833(19)$ Å, $\alpha = \gamma = 90.00$, $\beta = 96.442(8)^\circ$, $V = 4050.2(6)$ Å³, $Z = 8$, $d_{\text{calcd}} = 1.353$ g/cm³, $\mu = 0.112$ mm⁻¹, $F(000) = 1728$. Crystallographic data for compound **2a** (CCDC 924323) have been deposited at the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK.
- General procedures for the synthesis of compounds **3** and **4**. The corresponding chromane **2** (1.0 mmol) was refluxed in a mixture of EtOH (4 mL), H₂O (2 mL), and conc. HCl (0.2 mL) with vigorous stirring for 6 h. The reaction mixture was cooled to room temperature and the solid that formed was filtered, washed with H₂O (2 \times 1 mL), dried at 70 °C, and recrystallized from CH_2Cl_2 /hexane (1:2) (for **3**, a white powder) or MeOH (for **4**, colorless needles).
(2R*,3R*,4S*)-1-[6-Bromo-3-nitro-2-(trichloromethyl)-3,4-dihydro-2H-chromen-4-yl]-2,4-pentanedione (**3g**). Yield 0.30 g (63%), mp 140–141 °C. IR (KBr): 1715, 1626, 1601, 1552, 1480, 1412, 1366 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) enol **3g** (91%) δ 2.11 (s, 3H, Me), 2.57 (dd, 1H, CHH, $J = 16.4$, 10.7 Hz), 2.90 (dd, 1H, CHH, $J = 16.4$, 4.2 Hz), 3.87 (dd, 1H, H-4, $J = 10.7$, 4.2 Hz), 4.50 (br s, 1H, H-3), 5.53 (s, 1H, H-2), 5.71 (s, 1H, =CH), 7.01 (d, 1H, H-8, $J = 8.8$ Hz), 7.33 (d, 1H, H-5, $J = 2.2$ Hz), 7.38 (dd, 1H, H-7, $J = 8.8$, 2.2 Hz); diketo **3g** (9%) δ 2.76 (s, 3H, Me), 2.95–3.06 (m, 2H, CH₂), 3.63 (d, 1H, CHH, $J = 15.8$ Hz), 3.70 (d, 1H, CHH, $J = 15.8$ Hz), 3.96 (dd, 1H, H-4, $J = 8.2$, 5.5 Hz), 4.51 (br s, 1H, H-3), 5.61 (d, 1H, H-2, $J = 1.5$ Hz), 7.00 (d, 1H, H-8, $J = 8.9$ Hz), 7.28 (d, 1H, H-5, $J = 2.3$ Hz), 7.33 (dd, 1H, H-7, $J = 8.9$, 2.3 Hz); ^{13}C NMR (126 MHz, CDCl_3) enol **3g** δ 24.2, 36.9, 45.6, 78.8, 80.5, 95.2, 100.6, 115.4, 119.1, 122.9, 131.3, 131.8, 151.8, 189.9, 190.1; diketo **3g** δ 31.1, 35.3, 50.1, 56.9, 78.7, 80.6, 100.5, 115.3, 119.2, 122.8, 131.1, 131.7, 151.9, 200.2, 202.3. Anal. calcd for $\text{C}_{15}\text{H}_{13}\text{BrCl}_3\text{NO}_5$: C, 38.05; H, 2.77; N, 2.96. Found: C, 37.75; H, 2.59; N, 2.95.
1-[9-Bromo-2-hydroxy-5-(trifluoromethyl)-5H-chromeno[3,4-b]pyridin-3-yl]ethanone (**4c**). Yield 0.18 g (46%), mp 189–190 °C. IR (KBr): 1657, 1601, 1484, 1399, 1264, 1221 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 2.76 (s, 3H, Me), 5.60 (q, 1H, H-5, $J = 7.1$ Hz), 7.00 (d, 1H, H-7, $J = 8.8$ Hz), 7.47 (dd, 1H, H-8, $J = 8.8$, 2.1 Hz), 7.58 (s, 1H, H-1), 7.79 (d, 1H, H-10, $J = 2.1$ Hz), 11.93 (s, 1H, OH); ^{13}C NMR (126 MHz, CDCl_3) δ 25.7 (Me), 75.2 (q, C-5, $^2J_{\text{CF}} = 31.6$ Hz), 115.8 (C-9), 118.7 (C-1), 119.6 (C-7), 119.8 (C-10a), 123.0 (q, CF_3 , $^1J_{\text{CF}} = 286.9$ Hz), 127.1 (C-10), 130.9 (C-10b), 134.8 (C-4a), 135.1 (C-8), 135.6 (C-3), 151.6 (C-6a), 159.6 (C-2), 206.5 (C=O); ^{19}F NMR (471 MHz, CDCl_3) δ 85.2 (d, CF_3 , $J_{\text{FH}} = 7.1$ Hz); ^{15}N NMR (51 MHz, CDCl_3) δ 316.9; MS (EI): m/z 389 [⁸¹Br, M+1]⁺ (32), 387 [⁷⁹Br, M+1]⁺ (32), 320 [M+1– CF_3]⁺ (32), 138 [M–1– CF_3]⁺ (100), 290 [M+1– CF_3 –CO]⁺ (18), 196 (17), 169 (13), 140 (19), 113 (24), 63 (21), 43 [Ac]⁺ (7.9). Anal. calcd for $\text{C}_{15}\text{H}_9\text{BrF}_3\text{NO}_3$: C, 46.42; H, 2.34; N, 3.61. Found: C, 46.18; H, 2.48; N, 3.55.
- (a) Banini, S. R.; Turner, M. R.; Cummings, M. M.; Söderberg, B. C. G. *Tetrahedron* **2011**, *67*, 3603; (b) Madapa, S.; Singh, V.; Batra, S. *Tetrahedron* **2006**, *62*, 8740; (c) Lee, K. Y.; Kim, J. M.; Kim, J. N. *Tetrahedron* **2003**, *59*, 385; (d) Basavaiah, D.; Reddy, B. S.; Badsara, S. S. *Chem. Rev.* **2010**, *110*, 5447.
- Raczynska, E. D.; Kosińska, W.; Ośmiałowski, B.; Gawinecki, R. *Chem. Rev.* **2005**, *105*, 3561.
- Korotaev, V. Y.; Barkov, A. Y.; Slepukhin, P. A.; Kodess, M. I.; Sosnovskikh, V. Y. *Mendeleev Commun.* **2011**, *21*, 277.
- Levy, G.C.; Lichter, R.L. *Nitrogen-15 Nuclear Magnetic Resonance Spectroscopy*; John Wiley and Sons, 1979, p 80.
- Sheldrick, G. M. *Acta Crystallogr., Sect. A* **2008**, *64*, 112.